Aon Benfield
Analytics | Impact Forecasting

Hurricane Harvey
Event Recap Report

March 2018

Table of Contents

Executive Summary 3

Meteorological Recap 4

Storm Data 8

Impacted Areas & Effects 16

Texas 16

Louisiana 24

Energy Impacts 25

Transportation Impacts 28

Environmental Impacts 30

Comparison: HU Harvey vs. TS Allison (2001) 32

Impact Forecasting Reconnaissance Trip 35

Impact Forecasting: Modeled US Wind, Storm Surge, & Inland Flood Results 37

Hurricane Harvey ¶ V) L Q D Q F L D O , P S D F W 40

Appendix A: United States Hurricane Landfall Frequency 42

Appendix B: Costliest Atlantic Basin and United States Hurricanes 44

Appendix C: United States Emergency and Major Disaster Declarations 45

Appendix D: Glossary 46

Contact Information 49

Aon Bepof,(pi)-5epl

(8p)(p8(nc)-3apas)-(p)-10(r)-27py)18( )] TJ E 25 1 72188.6631.027 pypi0(w)1pr

Harvey maintained minimal tropical storm-force winds while meandering back into the Gulf of Mexico. The NHC only expected minimal (if any) intensification, primarily due to sea surface temperatures being much

Storm Data

Below is official storm data from the National Weather Service, Weather Prediction Center, and National Hurricane Center.

Wind & Central Pressure

Wind Gusts

Location	Wind Gust (mph)	Location	Wind Gust (mph)
Port Aransas, TX	132	Austwell, TX	80
Copano Village, TX	125	Magnolia Beach, TX	79
Lamar, TX	110	Edna, TX	73
Aransas Pass, TX	109	Chapman Ranch, TX	71

Select Rainfall Gauges





Rainfall Return Period

There is no simple way to provide a blanket rainfall return period (RP) for Hurricane Harvey. Based on typical rainfall patterns, there are some specific locations that end up receiving much higher – or lower – rainfall amounts. For Harvey, the extent of extreme rainfall was substantial. An unusually large and remarkable swath of minimal 40-inch (1,016-millimeter) rainfall extended from west of Houston to the Texas/Louisiana border. Due to this fact, preliminary rainfall return period analysis by NOAA's Hydrometeorological Design Studies Center indicates that an abnormally large swath of southeast Texas incurred rainfall RP rates minimally ranging from 500 to 1,000 years.

Given the volume of water, local infrastructure across southeast Texas was simply unable to handle such an enormous amount of rainfall in a short amount of time. This led to major water run-off that quickly accumulated across a very large area. With so much residential and commercial growth throughout this part of the state – combined with abundant concrete and poor absorbing clay soil – this only worsened the flood impact.

It is worth noting that Texas has incurred numerous extreme rainfall events during the past decade. In fact, Harvey became the sixth event to bring rainfall at a return period of at least 1,000 years since 2010 alone. The other events ocalone

River Gauge Height Map

Flooding stages across SE Texas and Houston area as of August 29, 11:00AM CDT (Data: NOAA, Background: OpenStreetMap)

Selected River Gauge Locations









Hydrographs for selected river gauges in Harris County (Source: NOAA)

Flood Damage

The majority of the damage inflicted by Harvey was as a result of catastrophic flooding inflicted by the combination of a large storm surge and prodigious amounts of rainfall as the storm pulled warm, moist air from the Gulf of Mexico over the Texas coast for days following its landfall. The NWS reported that conditions were “unprecedented” and issued a flash flood emergency for the entire Houston metro area.

River levels rose rapidly prompting widespread and devastating floods. Among the worst affected areas was Harris County, including the Houston metro area. By early on August 27, NWS meteorologists were already describing the floods as the “worst Houston had ever experienced” and official guidance from emergency management teams to people whose homes became inundated was to “get on the roof”. Another NWS official described the

By August 31, record-breaking river gauge levels were recorded in the Brazos River, Navidad River, San Bernard River, San Jacinto River, Trinity River, Clear Creek, Cypress Creek, Davidson Creek, Lake Creek, Menard Creek, Peach Creek, and Buffalo Bayou while the Colorado River, Guadalupe River, Lavaca River, Tres Palacios River, Bedias Creek, Caney Creek, Garcitas Creek, Sandies Creek, Sandy Creek, Spring Creek, Brays Bayou, White Oak Bayou, and Greens Bayou all reached major flood stage. Multiple other rivers, creeks, and bayous were in moderate or minor flood stages and many were still rising.

Emergency services were inundated by tens of thousands of calls for help and carried out thousands of

Summary of Property Damage in Texas

Single Family Homes & Businesses

County	Single Family Homes				Businesses	
	Affected	Minor Damage	Major Damage	Destroyed	Major Damage	Destroyed
Harris	51,011	38,640	24,790	845	359	0
Jefferson	30,193	15,405	16,131			

Public Buildings & Facilities

,1

County	Debris Removal	Police/EMS	Roads & Bridges	Water Facilities	Buildings	Public Utility Systems	Other	TOTAL
Harris	44,399,500	34,024,504	301,200	18,500	35,308,047	72,115,500	1,756,000	188,1

Louisiana

As Harvey edged northward through August 28-29, a federal disaster was declared for five “at-risk” Louisiana parishes including Beauregard, Calcasieu, Cameron, Jefferson Davis, and Vermillion in the southwest of the state. The Louisiana governor sought the addition of Allen, Acadia, Iberia, Natchitoches, Rapides, Sabine, and Vernon to the declaration on August 30. With rainfall topping 20 inches (508 millimeters) across southwestern sections of the state, this caused several rivers and streams to swell well above flood stage.

As of August 29, six refineries in the Corpus Christi area, seven in Houston/Galveston, and one in Beaumont/Port Arthur were shutdown. The combined capacity of these refineries was 3,268,449 barrels per day, equal to 33.7 percent of Gulf Coast refining capacity and 17.6 percent of total U.S. capacity. Furthermore, one refinery in Houston/Galveston, two in Beaumont/Port Arthur, and two in Lake Charles were operating at reduced rates. The following day, another Beaumont/Port Arthur refinery was added to the list of closures taking the affected capacity to 3,871,449 barrels per day or 39.9 percent of total Gulf Coast capacity: 20.9 percent of total U.S. capacity.

August 28, saw the first Corpus Christi refinery begin startup operations. Further startups were initiated on August 30 when six refineries with a capacity of 1,269,720 barrels per day (or 13.1 percent of Gulf Coast refining capacity) began the process. As of September 1, four refineries were operating at reduced capacities, six had initiated the startup process, and 10 remained shut down.

By September 4, eight refineries remained shut down with a combined capacity of 2,110,229 barrels per dayegainegainegain,equal 11 2.8[(13)4(.1)-8(pe)4(r)-3(c)-5(en)42(y)18(Tm [( )] TJ ET BT 020 0 1 87.984 516.91 Tm [(

B

Environmental Impacts

Coastal Erosion

Harvey was responsible for severe erosion all along the central and northern Texas coastline. Impacts

Comparison: HU Harvey vs.

Miscellaneous

NFIP Coverage

The graphic below shows NFIP coverage by county across the hardest-hit areas by Harvey. The data is based on statistics from NFIP and the U.S. Census Bureau at the time of landfall. In Harris County, TX – home to the city of Houston – just one out of six homes currently had active NFIP policies in place. The rates of coverage are most pronounced immediately along the coastline, but decline into the single-digits once getting further inland.

Impact Forecasting Reconnaissance Trip

Members of Aon Benfield's Catastrophe Management and Impact Forecasting teams traveled to Houston, Texas from September 19-21, 2017 for a three-day damage assessment following Hurricane Harvey's landfall in late August. Several neighborhoods and parts of the metro region were visited: Bellaire, Dickinson, Pearland, Meyerland, South Houston, Katy (Cinco Ranch), Buffalo Bayou reservoir (Nottingham Forest neighborhood), and Sienna Plantation.

Among the most notable findings:

1) There was a significant difference in areas with a mixture of homes built Pre-FIRM (Flood Insurance Rate Map) and those with newer construction in modern FEMA map zones. Homes which were elevated to meet current mandates often escaped without major damage. However, Pre-FIRM homes that were grandfathered with previous code standards often incurred entire loss to their first floor contents.

Home without any foundation elevation (Source: Impact Forecasting)

Home built with 12-18" elevation (Source: Impact Forecasting)

2) The height and longevity of flood waters across many neighborhoods and commercial regions led to significant molding and weakening of dry wall. The combination of tropical heat and humidity in Houston additionally worsened structures that did not have much external circulation. This often led to a loss of contents on the second story.

First floor remnants following 2' of water (Source: Impact Forecasting)

Molding on walls (Source: Impact Forecasting)

3) Significant flooding occurred in areas even where local officials had properly planned for at least a 100-year flood event. Given the extensive volume of new construction and concrete throughout Houston metro, the current infrastructure was not prepared to handle more than 60 inches of rainfall. This led to major flash flooding and inundation. The dominant soil type (clay) and excessive amount of concrete additionally provided further risk for flash flooding.

4) Dozens of tornado touchdowns during Hurricane Harvey led to roof damage in multiple neighborhoods. In some instances, roof damage was severe enough to allow for indoor water intrusion that led to an entire loss of indoor contents.

5) Water heights reached up to 8 feet in numerous neighborhoods, making ground rescue impossible. This led to some residents needing to be rescued from their roofs via helicopter after holes were punctured through attics.

Impact Forecasting: Modeled U.S.

As a reminder, data used for storm tide is water height above the North American Vertical Datum of 1988 (NAVD88). Note that this is different from inundation, which is the measured water height above ground level after high tide that occurs as a result of the storm tide. To see a full glossary of tropical cyclone-related terms see Appendix D.

Impact Forecasting Modeled Loss Results

Based on the modeled results of Impact Forecasting's wind, storm surge, and inland flood models from its initial near real-time runs in September 2017, it was determined that "insurable losses" resulting from Hurricane Harvey in the United States would range between USD52 and 69 billion . Please note that *"insurable losses" does not indicate th*e total economic loss estimate, nor the insured loss estimate. This only includes the value of properties (homes, businesses, vehicles) that are potentially insured by the private market

Financial Impact

Hurricane Harvey left a catastrophic path of devastation across a wide swath of southeast Texas and portions of extreme southwest Louisiana and is expected to have an overall economic impact in the vicinity of USD100

Appendix A

United States Hurricane Landfall Return Periods

Below are maps from the National Hurricane Center (NHC) that provide tropical cyclone return periods for areas along the United States coastline located south of the North Carolina border.

The first map shows regions at greatest risk of hurricanes tracking within 50 nautical miles (57 miles (93 kilometers)) of a particular location along the coastline. Historical data shows that the southern tip of Florida, Louisiana, and the outer banks of North Carolina are areas that typically record a hurricane every five to seven years.

The second map shows regions at greatest risk of major hurricanes within 50 nautical miles (57 miles (93 kilometers)) of a particular location along the coastline. Historical data shows, unsurprisingly, that there are longer return periods for stronger storms. Areas with greatest historical frequency include south Florida, the Florida panhandle, Louisiana, South Carolina, and North Carolina Harvey's first official landfall came in Aransas County, Texas, which has a return period for such an event every 33 - 40 years.

In terms of tropical cyclone development, the official Atlantic Hurricane season runs from June 1 to November 30 – though tropical cyclones have been known to develop during any month. As seen in the graphics below, the most likely months for cyclogenesis are during the peak months of August, September, and October. During these months, atmospheric and oceanic conditions are climatologically the most conducive for cyclones being influenced and/or steered by a strong Atlantic ridge of high pressure and/or by the advancement of strong troughs that dig into the central and eastern United States.

The caveat to cyclogenesis is, of course, the eventual track of storms that do develop. While g(s)] TJ S, tm ET BT 1 0 0

2058.8 . og(s)d cces rea5(op)4(.-10(i)522)-10(a s)-2-5(i)6(s)t d c 5(e to)4(g A)-4

Appendix B

Costliest Atlantic Basin and United States Hurricanes

Atlantic Basin

Rank	Storm Name	Affected Locations	Economic Loss [1] (2018 USD)	Insured Loss [2] (2018 USD)

Appendix C

United States Emergency Declarations

Texas

Emergency Declaration made on 08/23/2017.

Counties:

Aransas, Austin, Bastrop, Bee, Bexar, Brazoria, Burleson, Calhoun, Chambers, Colorado, Comal, Dallas, DeWitt, Fayette, Fort Bend, Galveston, Goliad, Gonzales, Grimes, Guadalupe, Hardin, Harris, Jackson, Jasper, Jefferson, Jim Wells, Karnes, Kleberg, Lavaca, Lee, Liberty, Madison, Matagorda, Milam, Montgomery, Newton, Nueces, Orange, Polk, Refugio, Sabine, San Augustine, San Jacinto, San Patricio, Tarrant, Travis, Tyler, Victoria, Walker, Waller, Washington, Wharton

Louisiana

Emergency Declaration made on 08/28/2017.

Parishes: Acadia, Allen, Ascension, Assumption, Avoyelles, Beauregard, Bienville, Bossier, Caddo, Calcasieu, Caldwell, Cameron, Catahoula, Claiborne, Concordia, DeSoto, East Baton Rouge, East Carroll, East Feliciana, Evangeline, Franklin, Grant, Iberia, Iberville, Jackson, Jefferson, Jefferson Davis, La Salle, Lafayette, Lafourche, Lincoln, Livingston, Madison, Morehouse, Natchitoches, Orleans, Ouachita, Plaquemines, Pointe Coupee, Rapides, Red River, Richland, Sabine, St. Bernard, St. Charles, St. Helena, St. James, St. John the Baptist, St. Landry, St. Martin, St. Mary, St. Tammany, Tangipahoa, Tensas, Terrebonne, Union, Vermilion, Vernon, Washington, Webster, West Baton Rouge, West Carroll, West Feliciana, and Winn.

United States Major Disaster Declarations

Texas

Major Disaster Declaration made on 08/25/2017.

Counties: Aransas, Austin, Bastrop, Bee, Brazoria, Calhoun, Chambers, Colorado, DeWitt, Fayette, Fort Bend, Galveston, Goliad, Gonzales, Grimes, Hardin, Harris, Jackson, Jasper, Jefferson, Karnes, Kleberg,

Appendix D

Glossary: Terms as defined by the National Hurricane Center

Advisory:
Official information issued by tropical cyclone

Contact Information

Adam Podlaha
Head of Impact Forecasting
Aon Benfield Analytics
Impact Forecasting
+44.20.7522.3820
adam.podlaha@aonbenfield.com

Siamak Daneshvaran
Head of Research & Development
Aon Benfield Analytics
Impact Forecasting
+1.312.381.5886
siamak.daneshvaran@aonbenfield.com

Anwesha Bhattacharya
Senior Analyst
Aon Benfield Analytics
Impact Forecasting
+ 91.080.6621.8575
anwesha.bhattec148(t)-1(@a)-6anowi-10(i)5(el)-4ld.com

sit3vw99(nf)-8(h)-9(n)-9w7(i)8-6(on)4(b)-9(en)4(f)-10(i)5(el)-4lopom